

Pedagogical formal and fonctionnal systems





Vincent Demange

CPR, Cédric, Cnam

12/06/2013

Introduction to formal pedagogy

Published works

-  L. Colson and D. Michel. *Pedagogical natural deduction systems : the propositional case*. JUCS, 13(10) :1396–1410, 2007.
-  L. Colson and D. Michel. *Pedagogical Second-Order Propositional Calculi*. JLC, 18(4) :669–695, 2008.
-  L. Colson and D. Michel. *Pedagogical second-order λ -calculus*. TCS, 410 :4190–4203, 2009.
-  L. Colson and V. Demange. *Investigations on a Pedagogical Calculus of Constructions*. JUCS, to appear, 2013.

Introduction to formal pedagogy

Basics : the pedagogical constraint

Poincaré criterion

«A definition by postulate has value **only if** there exists an example.» [Henri Poincaré – Last Thoughts (1913)]

Good postulate

Let x be a natural number such that $x^2 - 1 = 0$ holds.

▷ $x := 1$ suits

Bad postulate

Let y be a natural number verifying $y^2 + 1 = 0$.

▷ no example

Called “pedagogical” because of usual teaching practice

Outline of the investigation

1. Summary of the previous works
2. First attempts toward a pedagogical CC
3. A formal definition and some examples
4. Toward a Pedagogical Calculus of Constructions

Summary of the previous works

The formal pedagogical constraint

Poincaré criterion (informally)

«A definition by postulate has value **only if** there exists an example.» [Henri Poincaré – Last Thoughts (1913)]

Poincaré criterion (formally)

If \top is a type and o a term of type \top :

- ▶ $\Gamma := \{f : (\alpha \rightarrow \alpha) \rightarrow \alpha, g : \top \rightarrow \beta\}$ ▷ defines α and β
- ▶ $\sigma := [\alpha \mapsto \top, \beta \mapsto \top]$ exemplifies Γ :

$$\begin{aligned} &\vdash \lambda h^{\top \rightarrow \top}. o : (\top \rightarrow \top) \rightarrow \top \\ &\vdash \lambda x^{\top}. x : \top \rightarrow \top \end{aligned}$$

$\vdash \sigma \cdot \Gamma$ abbreviates that σ exemplifies Γ

Summary of the previous works

Simply typed λ -calculus

Morphology $\alpha \mid A \rightarrow B$

Syntax $x \mid \lambda x^A.u \mid u v$

$$\frac{x : F \in \Gamma}{\Gamma \vdash x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \vdash u : B}{\Gamma \vdash \lambda x^A.u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \vdash u : A \rightarrow B \quad \Gamma \vdash v : A}{\Gamma \vdash u v : B} \text{ (app)}$$

Summary of the previous works

Pedagogical simply typed λ -calculus [Colson and Michel (2007)]

Morphology $\alpha \mid A \rightarrow B$

Syntax $x \mid \lambda x^A.u \mid u v$

$$\frac{x : F \in \Gamma \quad \vdash \sigma \cdot \Gamma}{\Gamma \vdash x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \vdash u : B}{\Gamma \vdash \lambda x^A.u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \vdash u : A \rightarrow B \quad \Gamma \vdash v : A}{\Gamma \vdash u v : B} \text{ (app)}$$

Summary of the previous works

Pedagogical simply typed λ -calculus [Colson and Michel (2007)]

Morphology $\alpha \mid A \rightarrow B$

Syntax $x \mid \lambda x^A.u \mid u v$

No starting rule

$$\frac{x : F \in \Gamma \quad \vdash \sigma \cdot \Gamma}{\Gamma \vdash x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \vdash u : B}{\Gamma \vdash \lambda x^A.u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \vdash u : A \rightarrow B \quad \Gamma \vdash v : A}{\Gamma \vdash u v : B} \text{ (app)}$$

Summary of the previous works

Pedagogical simply typed λ -calculus [Colson and Michel (2007)]

Morphology $\alpha \mid A \rightarrow B \mid \top$

Syntax $x \mid \lambda x^A.u \mid u v \mid o$

$$\frac{\vdash \sigma \cdot \Gamma}{\Gamma \vdash o : \top} \text{ (ax)}$$

$$\frac{x : F \in \Gamma \quad \vdash \sigma \cdot \Gamma}{\Gamma \vdash x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \vdash u : B}{\Gamma \vdash \lambda x^A.u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \vdash u : A \rightarrow B \quad \Gamma \vdash v : A}{\Gamma \vdash u v : B} \text{ (app)}$$

Summary of the previous works

Pedagogical simply typed λ -calculus [Colson and Michel (2007)]

Morphology $\alpha \mid A \rightarrow B \mid \top$

Syntax $x \mid \lambda x^A.u \mid u v \mid o$

$$\frac{\vdash \sigma \cdot \Gamma}{\Gamma \vdash o : \top} \text{ (ax)}$$

$$\frac{x : F \in \Gamma \quad \vdash \sigma \cdot \Gamma}{\Gamma \vdash x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \vdash u : B}{\Gamma \vdash \lambda x^A.u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \vdash u : A \rightarrow B \quad \Gamma \vdash v : A}{\Gamma \vdash u v : B} \text{ (app)}$$

Results

- ▶ all formulas exemplified by \top
- ▶ initial and pedagogical systems (syntactically) equivalents

Summary of the previous works

System F [Girard (1972), Reynolds (1974)]

Morphology $\alpha \mid A \rightarrow B \mid \forall \alpha. A$

Syntax $x \mid \lambda x^A. u \mid u v \mid \Lambda \alpha. u$

$$\frac{x : F \in \Gamma}{\Gamma \vdash^f x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \vdash^f u : B}{\Gamma \vdash^f \lambda x^A. u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \vdash^f u : A \rightarrow B \quad \Gamma \vdash^f v : A}{\Gamma \vdash^f u v : B} \text{ (app)}$$

$$\frac{\Gamma \vdash^f u : B \quad \alpha \notin \mathcal{V}(\Gamma)}{\Gamma \vdash^f \Lambda \alpha. u : \forall \alpha. B} \text{ (Abs)}$$

$$\frac{\Gamma \vdash^f u : \forall \alpha. B}{\Gamma \vdash^f u V : B[\alpha \leftarrow V]} \text{ (App)}$$

Summary of the previous works

Weakly pedagogical System F [Colson and Michel (2008)]

Morphology $\alpha \mid A \rightarrow B \mid \forall \alpha. A \mid \top$

Syntax $x \mid \lambda x^A. u \mid u v \mid \Lambda \alpha. u \mid o$

$$\frac{\text{fw } \sigma \cdot \Gamma}{\Gamma \text{fw } o : \top} \text{ (ax)}$$

$$\frac{x : F \in \Gamma \quad \text{fw } \sigma \cdot \Gamma}{\Gamma \text{fw } x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \text{fw } u : B}{\Gamma \text{fw } \lambda x^A. u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \text{fw } u : A \rightarrow B \quad \Gamma \text{fw } v : A}{\Gamma \text{fw } u v : B} \text{ (app)}$$

$$\frac{\Gamma \text{fw } u : B \quad \alpha \notin \mathcal{V}(\Gamma)}{\Gamma \text{fw } \Lambda \alpha. u : \forall \alpha. B} \text{ (Abs)}$$

$$\frac{\Gamma \text{fw } u : \forall \alpha. B}{\Gamma \text{fw } u V : B[\alpha \leftarrow V]} \text{ (App)}$$

Summary of the previous works

Weakly pedagogical System F [Colson and Michel (2008)]

Morphology $\alpha \mid A \rightarrow B \mid \forall \alpha. A \mid \top$

Syntax $x \mid \lambda x^A. u \mid u v \mid \Lambda \alpha. u \mid o$

$$\frac{\Gamma \vdash^{\text{fw}} \sigma \cdot \Gamma}{\Gamma \vdash^{\text{fw}} o : \top} \text{ (ax)}$$

$$\frac{x : F \in \Gamma \quad \Gamma \vdash^{\text{fw}} \sigma \cdot \Gamma}{\Gamma \vdash^{\text{fw}} x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \vdash^{\text{fw}} u : B}{\Gamma \vdash^{\text{fw}} \lambda x^A. u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \vdash^{\text{fw}} u : A \rightarrow B \quad \Gamma \vdash^{\text{fw}} v : A}{\Gamma \vdash^{\text{fw}} u v : B} \text{ (app)}$$

$$\frac{\Gamma \vdash^{\text{fw}} u : B \quad \alpha \notin \mathcal{V}(\Gamma)}{\Gamma \vdash^{\text{fw}} \Lambda \alpha. u : \forall \alpha. B} \text{ (Abs)}$$

$$\frac{\Gamma \vdash^{\text{fw}} u : \forall \alpha. B}{\Gamma \vdash^{\text{fw}} u V : B[\alpha \leftarrow V]} \text{ (App)}$$

No subject reduction property :

$\Gamma \vdash^{\text{fw}} (\Lambda \alpha. \lambda x^\alpha. x) \perp : \perp \rightarrow \perp$ but

$\not\vdash^{\text{fw}} \lambda x^\perp. x : \perp \rightarrow \perp$

$\triangleright \perp := \forall \alpha. \alpha$

Summary of the previous works

Pedagogical System F [Colson and Michel (2009)]

Pre-morphology $\alpha \mid A \rightarrow B \mid \forall \alpha. A \mid \top$

Syntax $x \mid \lambda x^A. u \mid u v \mid \Lambda \alpha. u \mid o$

$$\frac{\text{fp } \sigma \cdot \Gamma}{\Gamma \text{fp } o : \top} \text{ (ax)}$$

$$\frac{x : F \in \Gamma \quad \text{fp } \sigma \cdot \Gamma}{\Gamma \text{fp } x : F} \text{ (var)}$$

$$\frac{\Gamma, x : A \text{fp } u : B}{\Gamma \text{fp } \lambda x^A. u : A \rightarrow B} \text{ (abs)}$$

$$\frac{\Gamma \text{fp } u : A \rightarrow B \quad \Gamma \text{fp } v : A}{\Gamma \text{fp } u v : B} \text{ (app)}$$

$$\frac{\Gamma \text{fp } u : B \quad \alpha \notin \mathcal{V}(\Gamma)}{\Gamma \text{fp } \Lambda \alpha. u : \forall \alpha. B} \text{ (Abs)}$$

$$\frac{\Gamma \text{fp } u : \forall \alpha. B \quad \text{fp } \sigma \cdot V}{\Gamma \text{fp } u V : B[\alpha \leftarrow V]} \text{ (App)}$$

Summary of the previous works

Pedagogical System F [Colson and Michel (2009)]

Main results

- ▶ every well-formed sub-type can be exemplified :

- ▶ Poincaré criterion always satisfied :

$$\Gamma \Vdash^{\text{fp}} u : A \Rightarrow \Vdash^{\text{fp}} \sigma \cdot \Gamma$$

- ▶ usefulness of functions :

$$\Gamma \Vdash^{\text{fp}} f : A \rightarrow B \text{ with } A \text{ closed} \Rightarrow A \text{ is inhabited}$$

- ▶ negationless second-order propositional calculus

- ▶ embeddings from System F to its pedagogical version :

- ▶ at logical side :

$$\exists t \Gamma \Vdash^{\text{f}} t : F \Leftrightarrow \exists t' \Gamma^\gamma \Vdash^{\text{fp}} t' : F^\gamma$$

where F^γ is F with occurrences of variables α replaced by $\alpha \vee \gamma$ (γ fresh)

- ▶ at computational side :

$$\Gamma \Vdash^{\text{f}} t : A \Rightarrow \Gamma^\delta \Vdash^{\text{fp}} \bar{t} : A^\delta$$

where A^δ is a *double-negation* version of A (\perp replaced by δ)
and \bar{t} is a CPS version of t (δ fresh)

Summary of the previous works

Pedagogical System F [Colson and Michel (2009)]

Main results

logic	computation
formula	datatype
proof	program
simplification	computation
proof checking	type checking
exemplifiability	utility

First attempts toward a pedagogical CC

Calculus of Constructions [Coquand and Huet (1985)]

Motivations

The Calculus of Constructions :

- ▶ uniform presentation of previous systems
- ▶ interdependence between morphology and syntax
 - ▷ reduction of constraints number
- ▶ first step towards pedagogical pure type systems (PTS)
 - ▷ especially Type : Type [Martin-Löf (1971)]

First attempts toward a pedagogical CC

Calculus of Constructions [Coquand and Huet (1985)]

$$\frac{}{\emptyset \text{ wf}^{\mathcal{C}}} \text{ (c-env}_1\text{)}$$

$$\frac{\Gamma \Vdash^{\mathcal{C}} A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}^{\mathcal{C}}} \text{ (c-env}_2\text{)}$$

$$\frac{\Gamma \text{ wf}^{\mathcal{C}}}{\Gamma \Vdash^{\mathcal{C}} \text{Prop} : \text{Type}} \text{ (c-ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{ wf}^{\mathcal{C}}}{\Gamma, x : A, \Gamma' \Vdash^{\mathcal{C}} x : A} \text{ (c-var)}$$

$$\frac{\Gamma, x : A \Vdash^{\mathcal{C}} u : B : \kappa}{\Gamma \Vdash^{\mathcal{C}} \lambda x^A. u : \forall x^A. B} \text{ (c-abs)}$$

$$\frac{\Gamma \Vdash^{\mathcal{C}} u : \forall x^A. B \quad \Gamma \Vdash^{\mathcal{C}} v : A}{\Gamma \Vdash^{\mathcal{C}} u v : B[x \leftarrow v]} \text{ (c-app)}$$

$$\frac{\Gamma, x : A \Vdash^{\mathcal{C}} B : \kappa}{\Gamma \Vdash^{\mathcal{C}} \forall x^A. B : \kappa} \text{ (c-prod)}$$

$$\frac{\Gamma \Vdash^{\mathcal{C}} u : A \quad A \simeq A' \quad \Gamma \Vdash^{\mathcal{C}} A' : \kappa}{\Gamma \Vdash^{\mathcal{C}} u : A'} \text{ (c-conv)}$$

κ denotes Prop or Type

First attempts toward a pedagogical CC

Naive transposition

Formal Poincaré criterion

$\Gamma \equiv x_1 : A_1, \dots, x_n : A_n$ can be exemplified by terms t_1, \dots, t_n if

$$\begin{array}{c} \vdash t_1 : A_1 \\ \vdash t_2 : A_2[x_1 \leftarrow t_1] \\ \vdots \\ \vdash t_n : A_n[x_1, \dots, x_{n-1} \leftarrow t_1, \dots, t_{n-1}] \end{array}$$

Abbreviation : $\vdash \sigma \cdot \Gamma$ with $\sigma := [x_1 \mapsto t_1, \dots, x_n \mapsto t_n]$

First idea

Substitutes the well-formedness of an environment by its exemplifiability

$\triangleright \Gamma \text{ wf}$ replaced by $\vdash \sigma \cdot \Gamma$

First attempts toward a pedagogical CC

Naive transposition

$$\frac{\text{⊢}^n \sigma \cdot \Gamma}{\Gamma \text{⊢}^n \circ : \top : \text{Prop} : \text{Type}} \quad (\text{n-ax}) \qquad \frac{\text{⊢}^n \sigma \cdot (\Gamma, x : A, \Gamma')}{\Gamma, x : A, \Gamma' \text{⊢}^n x : A} \quad (\text{n-var})$$

But

▶ “exemplifiable” does not imply “well-formed” :

▶ $x_1 : \text{Type}$

▷ $\sigma := [x_1 \mapsto \text{Prop}]$

▶ $x_1 : \text{Prop}, x_2 : (\lambda H^{\top \rightarrow x_1}. \top) (\lambda y^{\top}. y)$

▷ $\sigma := [x_1 \mapsto \top; x_2 \mapsto \circ]$

▶ etc.

▶ no subject reduction

▷ $\perp \rightarrow \perp$ inhabited

First attempts toward a pedagogical CC

Calculus of Constructions [Coquand and Huet (1985)]

$$\frac{}{\emptyset \text{wf}^{\mathcal{C}}} \text{ (c-env}_1)$$

$$\frac{\Gamma \Vdash^{\mathcal{C}} A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{wf}^{\mathcal{C}}} \text{ (c-env}_2)$$

$$\frac{\Gamma \text{wf}^{\mathcal{C}}}{\Gamma \Vdash^{\mathcal{C}} \text{Prop} : \text{Type}} \text{ (c-ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{wf}^{\mathcal{C}}}{\Gamma, x : A, \Gamma' \Vdash^{\mathcal{C}} x : A} \text{ (c-var)}$$

$$\frac{\Gamma, x : A \Vdash^{\mathcal{C}} u : B : \kappa}{\Gamma \Vdash^{\mathcal{C}} \lambda x^A. u : \forall x^A. B} \text{ (c-abs)}$$

$$\frac{\Gamma \Vdash^{\mathcal{C}} u : \forall x^A. B \quad \Gamma \Vdash^{\mathcal{C}} v : A}{\Gamma \Vdash^{\mathcal{C}} u \ v : B[x \leftarrow v]} \text{ (c-app)}$$

$$\frac{\Gamma, x : A \Vdash^{\mathcal{C}} B : \kappa}{\Gamma \Vdash^{\mathcal{C}} \forall x^A. B : \kappa} \text{ (c-prod)}$$

$$\frac{\Gamma \Vdash^{\mathcal{C}} u : A \quad A \simeq A' \quad \Gamma \Vdash^{\mathcal{C}} A' : \kappa}{\Gamma \Vdash^{\mathcal{C}} u : A'} \text{ (c-conv)}$$

κ denotes Prop or Type

First attempts toward a pedagogical CC

Calculus of Constructions [Coquand and Huet (1985)]

$$\frac{}{\emptyset \text{ wf}^{\text{C}}} \text{ (c-env}_1\text{)}$$

$$\frac{\Gamma \Vdash^{\text{C}} A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}^{\text{C}}} \text{ (c-env}_2\text{)}$$

$$\frac{\Gamma \text{ wf}^{\text{C}}}{\Gamma \Vdash^{\text{C}} \text{Prop} : \text{Type}} \text{ (c-ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{ wf}^{\text{C}}}{\Gamma, x : A, \Gamma' \Vdash^{\text{C}} x : A} \text{ (c-var)}$$

$$\frac{\Gamma, x : A \Vdash^{\text{C}} u : B : \kappa}{\Gamma \Vdash^{\text{C}} \lambda x^A. u : \forall x^A. B} \text{ (c-abs)}$$

$$\frac{\Gamma \Vdash^{\text{C}} u : \forall x^A. B \quad \Gamma \Vdash^{\text{C}} v : A}{\Gamma \Vdash^{\text{C}} u \ v : B[x \leftarrow v]} \text{ (c-app)}$$

$$\frac{\Gamma, x : A \Vdash^{\text{C}} B : \kappa}{\Gamma \Vdash^{\text{C}} \forall x^A. B : \kappa} \text{ (c-prod)}$$

$$\frac{\Gamma \Vdash^{\text{C}} u : A \quad A \simeq A' \quad \Gamma \Vdash^{\text{C}} A' : \kappa}{\Gamma \Vdash^{\text{C}} u : A'} \text{ (c-conv)}$$

(c-prod) responsible of vacuity

κ denotes Prop or Type

First attempts toward a pedagogical CC

Poincaréan Calculus of Constructions – CCr

Strategy change

- ▶ keep Γ wf judgements
- ▶ avoid empty types as soon as possible

Radical idea

Constrain (only) the type formation rule :

$$\frac{\Gamma, x : A \vdash^c B : \kappa}{\Gamma \vdash^c \forall x^A. B : \kappa} \text{ (c-prod)}$$

First attempts toward a pedagogical CC

Poincaréan Calculus of Constructions – CCr

Strategy change

- ▶ keep Γ wf judgements
- ▶ avoid empty types as soon as possible

Radical idea

Constrain (only) the type formation rule :

$$\frac{\Gamma, x : A \vdash^r u : B : \kappa}{\Gamma \vdash^r \forall x^A. B : \kappa} \text{ (r-prod)}$$

First attempts toward a pedagogical CC

Poincaréan Calculus of Constructions – CCr

Theorems

- ▶ respectful of the Poincaré criterion
 - ▷ relies on strong normalization of CC
- ▶ subject reduction property holds
 - ▷ Coq proof adapted from [Barras (1996)]
- ▶ usefulness of functions
 - ▷ $\vdash^r f : \forall x^A. B \Rightarrow \exists u \vdash^r u : A$
- ▶ terms of Gödel system T can be interpreted in CC_r
 - ▷ usual way : recursion from iteration (and trick for cartesian products)

But

- ▶ does not contain natively simply typed λ -calculus
 - ▷ $\nexists u \ A \ B \ C : \text{Prop} \vdash^r u : (A \rightarrow B) \rightarrow (B \rightarrow C) \rightarrow (A \rightarrow C)$
- ▶ does not prove symmetry of Leibniz equality
 - ▷ $\nexists u \ A : \text{Prop}, x \ y \ z : A \vdash^r u : x =_A y \rightarrow y =_A z \rightarrow x =_A z$

First attempts toward a pedagogical CC

Poincaréan Calculus of Constructions – CCr

Observation CCr seems too much constrained

Goal increase expressive power

Idea exemplifiable types should be usable

▷ similar to pedagogical system F

Definition : converse of the Poincaré criterion

CC_* , sub-system of CC, meets the *converse of the Poincaré criterion* if :

$$\vdash^* \sigma \cdot \Gamma \quad \Rightarrow \quad \Gamma \text{ wf}^*$$

First attempts toward a pedagogical CC

Poincaréan Calculus of Constructions – CCr

Observation CCr seems too much constrained

Goal increase expressive power

Idea exemplifiable types should be usable

▷ similar to pedagogical system F

Beware

Exemplifiable types need not be well-formed

Definition : converse of the Poincaré criterion

CC_* , sub-system of CC, meets the *converse of the Poincaré criterion* if :

$$\vdash^* \sigma \cdot \Gamma \text{ and } \Gamma \text{ wf}^c \Rightarrow \Gamma \text{ wf}^*$$

A formal definition and some examples

Pedagogical sub-system of the Calculus of Constructions

Definition : CC_{\star} pedagogical sub-system of CC

- ▶ CC_{\star} sub-system of CC

$$\triangleright \Gamma \vdash^{\star} u : A \Rightarrow \Gamma \vdash^{\mathbb{C}} u : A$$

- ▶ Subject reduction property holds for CC_{\star}

$$\triangleright \Gamma \vdash^{\star} u : A \text{ and } u \rightsquigarrow u' \Rightarrow \Gamma \vdash^{\star} u' : A$$

- ▶ CC_{\star} meets Poincaré criterion and its converse

$$\triangleright \Gamma \text{ wf}^{\star} \Leftrightarrow \vdash^{\star} \sigma \cdot \Gamma \text{ and } \Gamma \text{ wf}^{\mathbb{C}}$$

A formal definition and some examples

Calculus of Constructions [Coquand and Huet (1985)]

$$\frac{}{\emptyset \text{ wf}^c} \text{ (c-env}_1\text{)}$$

$$\frac{\Gamma \Vdash A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}^c} \text{ (c-env}_2\text{)}$$

$$\frac{\Gamma \text{ wf}^c}{\Gamma \Vdash \text{Prop} : \text{Type}} \text{ (c-ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{ wf}^c}{\Gamma, x : A, \Gamma' \Vdash x : A} \text{ (c-var)}$$

$$\frac{\Gamma, x : A \Vdash u : B : \kappa}{\Gamma \Vdash \lambda x^A. u : \forall x^A. B} \text{ (c-abs)}$$

$$\frac{\Gamma \Vdash u : \forall x^A. B \quad \Gamma \Vdash v : A}{\Gamma \Vdash u \ v : B[x \leftarrow v]} \text{ (c-app)}$$

$$\frac{\Gamma, x : A \Vdash B : \kappa}{\Gamma \Vdash \forall x^A. B : \kappa} \text{ (c-prod)}$$

$$\frac{\Gamma \Vdash u : A \quad A \simeq A' \quad \Gamma \Vdash A' : \kappa}{\Gamma \Vdash u : A'} \text{ (c-conv)}$$

κ denotes Prop or Type

A formal definition and some examples

Second order λ -calculus – λ^2

$$\frac{}{\emptyset \text{ wf}^2} \text{ (env}_1\text{)} \qquad \frac{\Gamma \models A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}^2} \text{ (env}_2\text{)}$$

$$\frac{\Gamma \text{ wf}^2}{\Gamma \models \text{Prop} : \text{Type}} \text{ (ax)} \qquad \frac{\Gamma, x : A, \Gamma' \text{ wf}^2}{\Gamma, x : A, \Gamma' \models x : A} \text{ (var)}$$

$$\frac{\Gamma, x : A \models u : B : \text{Prop}}{\Gamma \models \lambda x^A. u : \forall x^A. B} \text{ (abs)} \qquad \frac{\Gamma \models u : \forall x^A. B \quad \Gamma \models v : A}{\Gamma \models u \ v : B[x \leftarrow v]} \text{ (app)}$$

$$\frac{\Gamma, x : A \models B : \text{Prop}}{\Gamma \models \forall x^A. B : \text{Prop}} \text{ (prod)}$$

A formal definition and some examples

With explicit total exemplifications – λ_e^2

Goal

Obtain a pedagogical version of λ^2

▷ in the sense of the previous definition

First idea

Keep examples explicit : $\Gamma \vdash_{\sigma} u : A$ and $\Gamma \text{ wf}_{\sigma}$ where

- ▶ $\Gamma \equiv x_1 : A_1, \dots, x_n : A_n$
- ▶ $\sigma \equiv [x_1 \mapsto t_1, \dots, x_n \mapsto t_n]$ or in short $\sigma \equiv [t_1, \dots, t_n]$

A formal definition and some examples

Second order λ -calculus – λ^2

$$\frac{}{\emptyset \text{ wf}^2} \text{ (env}_1\text{)}$$

$$\frac{\Gamma \Vdash A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}^2} \text{ (env}_2\text{)}$$

$$\frac{\Gamma \text{ wf}^2}{\Gamma \Vdash \text{Prop} : \text{Type}} \text{ (ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{ wf}^2}{\Gamma, x : A, \Gamma' \Vdash x : A} \text{ (var)}$$

$$\frac{\Gamma, x : A \Vdash u : B : \text{Prop}}{\Gamma \Vdash \lambda x^A. u : \forall x^A. B} \text{ (abs)}$$

$$\frac{\Gamma \Vdash u : \forall x^A. B \quad \Gamma \Vdash v : A}{\Gamma \Vdash u v : B[x \leftarrow v]} \text{ (app)}$$

$$\frac{\Gamma, x : A \Vdash B : \text{Prop}}{\Gamma \Vdash \forall x^A. B : \text{Prop}} \text{ (prod)}$$

A formal definition and some examples

With explicit total exemplifications – λ_e^2

$$\frac{}{\emptyset \text{ wf}_{\emptyset}^{2e}} \text{ (e-env}_1\text{)}$$

$$\frac{\Gamma \Vdash_{\sigma}^{2e} A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}_{\sigma :: a}^{2e}} \text{ (e-env}_2\text{)}$$

$$\frac{\Gamma \text{ wf}_{\sigma}^{2e}}{\Gamma \Vdash_{\sigma}^{2e} \text{ Prop} : \text{Type}} \text{ (e-ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{ wf}_{\sigma}^{2e}}{\Gamma, x : A, \Gamma' \Vdash_{\sigma}^{2e} x : A} \text{ (e-var)}$$

$$\frac{\Gamma, x : A \Vdash_{\sigma :: a}^{2e} u : B : \text{Prop}}{\Gamma \Vdash_{\sigma}^{2e} \lambda x^A. u : \forall x^A. B} \text{ (e-abs)}$$

$$\frac{\Gamma \Vdash_{\sigma}^{2e} u : \forall x^A. B \quad \Gamma \Vdash_{\sigma}^{2e} v : A}{\Gamma \Vdash_{\sigma}^{2e} u v : B[x \leftarrow v]} \text{ (e-app)}$$

$$\frac{\Gamma, x : A \Vdash_{\sigma :: a}^{2e} B : \text{Prop}}{\Gamma \Vdash_{\sigma}^{2e} \forall x^A. B : \text{Prop}} \text{ (e-prod)}$$

A formal definition and some examples

With explicit total exemplifications – λ_e^2

$$\frac{\text{---}}{\emptyset \text{ wf}_{\emptyset}^{2e}} \text{ (e-env}_1\text{)} \qquad \frac{\Gamma \Vdash_{\sigma}^{2e} A : \kappa \quad \Vdash_{\emptyset}^{2e} a : \sigma(A) \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}_{\sigma :: a}^{2e}} \text{ (e-env}_2\text{)}$$

$$\frac{\Gamma \text{ wf}_{\sigma}^{2e}}{\Gamma \Vdash_{\sigma}^{2e} o : \mathbb{T} : \text{Prop} : \text{Type}} \text{ (e-ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{ wf}_{\sigma}^{2e}}{\Gamma, x : A, \Gamma' \Vdash_{\sigma}^{2e} x : A} \text{ (e-var)}$$

$$\frac{\Gamma, x : A \Vdash_{\sigma :: a}^{2e} u : B : \text{Prop}}{\Gamma \Vdash_{\sigma}^{2e} \lambda x^A. u : \forall x^A. B} \text{ (e-abs)}$$

$$\frac{\Gamma \Vdash_{\sigma}^{2e} u : \forall x^A. B \quad \Gamma \Vdash_{\sigma}^{2e} v : A}{\Gamma \Vdash_{\sigma}^{2e} u v : B[x \leftarrow v]} \text{ (e-app)}$$

$$\frac{\Gamma, x : A \Vdash_{\sigma :: a}^{2e} B : \text{Prop} \quad \Vdash_{\emptyset}^{2e} t : \sigma(\forall x^A. B)}{\Gamma \Vdash_{\sigma}^{2e} \forall x^A. B : \text{Prop}} \text{ (e-prod)}$$

A formal definition and some examples

With explicit total exemplifications – λ_e^2

Theorems

- ▶ λ_e^2 (*almost*) a pedagogical sub-system of CC
- ▶ examples contained as sub-derivations
 - ▷ $\Gamma \vdash_{\sigma}^{2e} A : \kappa \Rightarrow \exists a \vdash_{\emptyset}^{2e} a : \sigma(A)$ sub-derivation
- ▶ exchange of exemplifications :
 - ▷ $\Gamma \vdash_{\sigma}^{2e} w : C$ and $\Gamma \text{wf}_{\rho}^{2e} \Rightarrow \Gamma \vdash_{\rho}^{2e} w : C$

But

λ_e^2 not exactly sub-system of CC because :

- ▶ exemplifications are explicit
- ▶ addition of two symbols o and \top

A formal definition and some examples

With explicit total exemplifications – λ_e^2

Theorems

- ▶ λ_e^2 (*almost*) a pedagogical sub-system of CC
- ▶ examples contained as sub-derivations
 - ▷ $\Gamma \vdash_{\sigma}^{2e} A : \kappa \Rightarrow \exists a \vdash_{\emptyset}^{2e} a : \sigma(A)$ sub-derivation
- ▶ exchange of exemplifications :
 - ▷ $\Gamma \vdash_{\sigma}^{2e} w : C$ and $\Gamma \text{ wf}_{\rho}^{2e} \Rightarrow \Gamma \vdash_{\rho}^{2e} w : C$

But

λ_e^2 not exactly sub-system of CC because :

- ▶ exemplifications are explicit
- ▶ addition of two symbols o and \top

Second idea

Relax constraints on exemplifications by making them implicit

A formal definition and some examples

With explicit total exemplifications – λ_e^2

$$\frac{}{\emptyset \text{ wf}_{\emptyset}^{2e}} \text{ (e-env}_1) \qquad \frac{\Gamma \Vdash_{\sigma}^{2e} A : \kappa \quad \Vdash_{\emptyset}^{2e} a : \sigma(A) \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}_{\sigma}^{2e} : a} \text{ (e-env}_2)$$

$$\frac{\Gamma \text{ wf}_{\sigma}^{2e}}{\Gamma \Vdash_{\sigma}^{2e} \circ : \top : \text{Prop} : \text{Type}} \text{ (e-ax)} \qquad \frac{\Gamma, x : A, \Gamma' \text{ wf}_{\sigma}^{2e}}{\Gamma, x : A, \Gamma' \Vdash_{\sigma}^{2e} x : A} \text{ (e-var)}$$

$$\frac{\Gamma, x : A \Vdash_{\sigma}^{2e} : a \quad u : B : \text{Prop}}{\Gamma \Vdash_{\sigma}^{2e} \lambda x^A. u : \forall x^A. B} \text{ (e-abs)} \qquad \frac{\Gamma \Vdash_{\sigma}^{2e} u : \forall x^A. B \quad \Gamma \Vdash_{\sigma}^{2e} v : A}{\Gamma \Vdash_{\sigma}^{2e} u v : B[x \leftarrow v]} \text{ (e-app)}$$

$$\frac{\Gamma, x : A \Vdash_{\sigma}^{2e} : a \quad B : \text{Prop} \quad \Vdash_{\emptyset}^{2e} t : \sigma(\forall x^A. B)}{\Gamma \Vdash_{\sigma}^{2e} \forall x^A. B : \text{Prop}} \text{ (e-prod)}$$

A formal definition and some examples

With implicit total exemplifications – λ_t^2

$$\frac{}{\emptyset \text{ wf}^{2t}} \text{ (t-env}_1\text{)} \qquad \frac{\Gamma \models^{2t} A : \kappa \qquad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}^{2t}} \text{ (t-env}_2\text{)}$$

$$\frac{\Gamma \text{ wf}^{2t}}{\Gamma \models^{2t} \circ : \top : \text{Prop} : \text{Type}} \text{ (t-ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{ wf}^{2t}}{\Gamma, x : A, \Gamma' \models^{2t} x : A} \text{ (t-var)}$$

$$\frac{\Gamma, x : A \models^{2t} \quad u : B : \text{Prop}}{\Gamma \models^{2t} \lambda x^A. u : \forall x^A. B} \text{ (t-abs)}$$

$$\frac{\Gamma \models^{2t} u : \forall x^A. B \quad \Gamma \models^{2t} v : A}{\Gamma \models^{2t} u v : B[x \leftarrow v]} \text{ (t-app)}$$

$$\frac{\Gamma, x : A \models^{2t} \quad B : \text{Prop} \quad \sigma \text{ mot}_\Gamma \forall x^A. B}{\Gamma \models^{2t} \forall x^A. B : \text{Prop}} \text{ (t-prod)}$$

where $\sigma \text{ mot}_\Gamma C$ abbreviates :

- (a) σ exemplifies Γ , i.e. $\models^{2t} \sigma \cdot \Gamma$
- (b) and there is term t such that $\models^{2t} t : \sigma(C)$

A formal definition and some examples

With implicit total exemplifications – λ_t^2

Theorems

λ_t^2 equivalent to $\lambda_e^2 : \Gamma \Vdash^t t : A \Leftrightarrow \exists \sigma \Gamma \Vdash_\sigma^e t : A$

- ▶ λ_t^2 (*almost*) a pedagogical sub-system of CC

But

λ_t^2 not exactly sub-system of CC because :

- ▶ addition of two symbols o and \top

A formal definition and some examples

With implicit total exemplifications – λ_t^2

Theorems

λ_t^2 equivalent to $\lambda_e^2 : \Gamma \Vdash^t t : A \Leftrightarrow \exists \sigma \Gamma \Vdash_\sigma^e t : A$

- ▶ λ_t^2 (*almost*) a pedagogical sub-system of CC

But

λ_t^2 not exactly sub-system of CC because :

- ▶ addition of two symbols \circ and \top

Last idea

Have partial exemplifications to restore the CC_r 's behaviour :

$\Gamma \Vdash \text{Id} : \text{True} : \text{Prop}$

▷ with $\text{Id} := \lambda A^{\text{Prop}}. \lambda x^A. x$ and $\text{True} := \forall A^{\text{Prop}}. A \rightarrow A$

A formal definition and some examples

With implicit total exemplifications – λ_t^2

$$\begin{array}{c}
 \frac{}{\emptyset \text{ wf}^{2t}} \text{ (t-env}_1\text{)} \qquad \frac{\Gamma \vdash^{2t} A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}^{2t}} \text{ (t-env}_2\text{)} \\
 \\
 \frac{\Gamma \text{ wf}^{2t}}{\Gamma \vdash^{2t} \circ : \top : \text{Prop} : \text{Type}} \text{ (t-ax)} \qquad \frac{\Gamma, x : A, \Gamma' \text{ wf}^{2t}}{\Gamma, x : A, \Gamma' \vdash^{2t} x : A} \text{ (t-var)} \\
 \\
 \frac{\Gamma, x : A \vdash^{2t} u : B : \text{Prop}}{\Gamma \vdash^{2t} \lambda x^A. u : \forall x^A. B} \text{ (t-abs)} \qquad \frac{\Gamma \vdash^{2t} u : \forall x^A. B \quad \Gamma \vdash^{2t} v : A}{\Gamma \vdash^{2t} u v : B[x \leftarrow v]} \text{ (t-app)} \\
 \\
 \frac{\Gamma, x : A \vdash^{2t} B : \text{Prop} \quad \sigma \text{ mot}_\Gamma \forall x^A. B}{\Gamma \vdash^{2t} \forall x^A. B : \text{Prop}} \text{ (t-prod)}
 \end{array}$$

where $\sigma \text{ mot}_\Gamma C$ abbreviates :

- (a) σ exemplifies Γ , i.e. $\vdash^{2t} \sigma \cdot \Gamma$
- (b) and there is term t such that $\vdash^{2t} t : \sigma(C)$

A formal definition and some examples

With implicit partial exemplifications – λ_p^2

$$\begin{array}{c}
 \frac{}{\emptyset \text{ wf}^{2p}} \text{ (p-env}_1\text{)} \qquad \frac{\Gamma \Vdash^p A : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{ wf}^{2p}} \text{ (p-env}_2\text{)} \\
 \\
 \frac{\Gamma \text{ wf}^{2p}}{\Gamma \Vdash^p \text{ Prop} : \text{Type}} \text{ (p-ax)} \qquad \frac{\Gamma, x : A, \Gamma' \text{ wf}^{2p}}{\Gamma, x : A, \Gamma' \Vdash^p x : A} \text{ (p-var)} \\
 \\
 \frac{\Gamma, x : A \Vdash^p u : B : \text{Prop}}{\Gamma \Vdash^p \lambda x^A. u : \forall x^A. B} \text{ (p-abs)} \qquad \frac{\Gamma \Vdash^p u : \forall x^A. B \quad \Gamma \Vdash^p v : A}{\Gamma \Vdash^p u v : B[x \leftarrow v]} \text{ (p-app)} \\
 \\
 \frac{\Gamma, x : A \Vdash^p B : \text{Prop} \quad \sigma \widetilde{\text{mot}}_{\Gamma} \forall x^A. B}{\Gamma \Vdash^p \forall x^A. B : \text{Prop}} \text{ (p-prod)}
 \end{array}$$

where $\sigma \widetilde{\text{mot}}_{\Gamma} C$ abbreviates :

- (a) σ partially exemplifies Γ , i.e. $\text{dom}(\sigma) \subseteq \text{dom}(\Gamma)$
- (b) and there is a term t such that $\sigma(\Gamma) \Vdash^p t : \sigma(C)$

A formal definition and some examples

With implicit partial exemplifications – λ_p^2

Theorems

- ▶ λ_p^2 equivalent to $\lambda_t^2 : \Gamma \Vdash^p t : A \iff \Gamma \Vdash^t t : A$
 - ▷ exemplifications can be completed
- ▶ λ_p^2 is a **pedagogical sub-system of CC**
 - ▷ definition exemplified!

General theorems

- ▶ λ_e^2 , λ_t^2 and λ_p^2 are equivalents
- ▶ embeddings to and from pedagogical system F
- ▶ embedding from pedagogical λ^2

A formal definition and some examples

With implicit partial exemplifications – λ_p^2

But

Type checking not decidable in λ_e^2 , λ_t^2 nor λ_p^2 :

$$\begin{aligned} \Gamma \vdash_{\sigma}^{2e} u : A \text{ ? dec.} &\Rightarrow \Gamma' \vdash^{2t} v : B \text{ ? dec.} \\ &\Rightarrow \exists ?w \Delta \vdash^{fp} w : C \text{ dec.} \\ &\Rightarrow \exists ?w' \Delta' \vdash^{f} w' : D \text{ dec.} \end{aligned}$$

▷ $\exists ?w' \Delta' \vdash^{f} w' : D$ undecidable [Urzyczyn (1997)]

Idea

Annotate types with terms to ensure exemplification :

$$\frac{\Gamma_{\sigma} \vdash A_a : \kappa \quad x \notin \text{dom}(\Gamma)}{\Gamma_{\sigma}, x : A_a \text{ wf}} \text{ (env}_2) \qquad \frac{\Gamma_{\sigma}, x : A_a \vdash B_b : \text{Prop} \quad \vdash t : \sigma(\forall x^{A_a}. B_b)}{\Gamma_{\sigma} \vdash (\forall x^{A_a}. B_b)_t : \text{Prop}} \text{ (prod)}$$

where $\Gamma_{\sigma} \equiv x_1 : A_{1_{a_1}}, \dots, x_n : A_{n_{a_n}}$

Toward a Pedagogical Calculus of Constructions

Higher order

Goal

Obtain a Pedagogical λ^ω

New objects

- ▶ predicates/propositional functions

$$\triangleright \lambda A^{\text{Prop}}.\lambda B^{\text{Prop} \rightarrow \text{Prop}}.B A$$

- ▶ postulated into environments

$$\triangleright f : \text{Prop} \rightarrow (\text{Prop} \rightarrow \text{Prop}) \rightarrow \text{Prop}$$

⇒ notion of predicate exemplification needed

Idea

Exemplifiable predicate \Leftrightarrow useful propositional function

Toward a Pedagogical Calculus of Constructions

Higher order

Exemplifiable predicate (formally)

$P : \mathcal{O}_1 \rightarrow \dots \rightarrow \mathcal{O}_n \rightarrow \text{Prop}$ exemplifiable if :

- ▶ can be completely applied
 - ▶ reducible to an exemplifiable type
- ▷ there are $u_i : \mathcal{O}_i$
- ▷ $P \vec{u} \rightsquigarrow^* R$ and $\vdash \sigma \cdot R$

Abbreviated : $\sigma \text{ mot}^{\mathcal{O}_1 \rightarrow \dots \rightarrow \mathcal{O}_n \rightarrow \text{Prop}}(P)$

Exemplifiable predicates

$P := \lambda A^{\text{Prop}}. A \rightarrow A$

$Q := \lambda A^{\text{Prop}}. \lambda B^{\text{Prop} \rightarrow \text{Prop}}. B A$

▷ $P \text{ T} \rightsquigarrow^* \text{T} \rightarrow \text{T}$

▷ $Q \text{ T} (\lambda C^{\text{Prop}}. \text{T}) \rightsquigarrow^* \text{T}$

Non exemplifiable predicate

$S := \lambda A^{\text{Prop}}. \forall B^{\text{Prop}}. A \leftrightarrow B$

$V := \lambda A^{\text{Prop}}. \forall B^{\text{Prop}}. A \rightarrow B$

▷ hopefully no arguments

Toward a Pedagogical Calculus of Constructions

Pedagogical higher-order λ -calculus – λ_e^ω

Beware interaction of exemplifiable predicates

► $Id := \lambda A^{\text{Prop}}.A$ and $Q := \lambda R^{\text{Prop} \rightarrow \text{Prop}}.\forall F^{\text{Prop}}.R F$

▷ but $Q Id \rightsquigarrow^* \perp$

Necessary constraint

$$\frac{\Gamma \vdash_\sigma u : A \rightarrow B : \text{Type} \quad \Gamma \vdash_\sigma v : A \quad \sigma \text{ mot}^B(u v)}{\Gamma \vdash_\sigma u v : B} \text{ (we-app}\square\text{)}$$

Toward a Pedagogical Calculus of Constructions

Pedagogical higher-order λ -calculus – λ_e^ω

Beware interaction of exemplifiable predicates

- ▶ $Id := \lambda A^{\text{Prop}}.A$ and $Q := \lambda R^{\text{Prop} \rightarrow \text{Prop}}.\forall F^{\text{Prop}}.R F$
▷ but $Q Id \rightsquigarrow^* \perp$
- ▶ $S := \lambda A^{\text{Prop} \rightarrow \text{Prop}}.\lambda H^{\forall B^{\text{Prop}}.A B}.H$
▷ but $S Id \rightsquigarrow^* \lambda H^\perp.H$

Necessary constraint

$$\frac{\Gamma \vdash_\sigma u : \forall x^A.B : \text{Prop} \quad \Gamma \vdash_\sigma v : A \quad \sigma \text{ mot}^{\text{Prop}}(B[x \leftarrow v])}{\Gamma \vdash_\sigma u v : B[x \leftarrow v]} \text{ (we-app}_*\text{)}$$

Toward a Pedagogical Calculus of Constructions

Pedagogical higher-order λ -calculus – λ_e^ω

Beware interaction of exemplifiable predicates

- ▶ $Id := \lambda A^{\text{Prop}}.A$ and $Q := \lambda R^{\text{Prop} \rightarrow \text{Prop}}.\forall F^{\text{Prop}}.R F$
▷ but $Q Id \rightsquigarrow^* \perp$
- ▶ $S := \lambda A^{\text{Prop} \rightarrow \text{Prop}}.\lambda H^{\forall B^{\text{Prop}}.A} B.H$
▷ but $S Id \rightsquigarrow^* \lambda H^\perp.H$

Necessary constraint

$$\frac{\Gamma \vdash_\sigma u : \forall x^A.B : \text{Prop} \quad \Gamma \vdash_\sigma v : A \quad \sigma \text{ mot}^{\text{Prop}}(B[x \leftarrow v])}{\Gamma \vdash_\sigma u v : B[x \leftarrow v]} \text{ (we-app}_*\text{)}$$

Strong : only normal form examples ?

Toward a Pedagogical Calculus of Constructions

Pedagogical higher-order λ -calculus – λ_e^ω

Theorems

- ▶ λ_e^ω satisfies Poincaré criterion
- ▶ λ_e^ω satisfies the converse Poincaré criterion
- ▶ **subject reduction still conjectured**

▷ usual substitution lemma invalid :

$$A : \star \rightarrow \star \vdash_{\lambda Z^*, \top} \lambda H^{\forall B^*. A B}. H : (\forall B^*. A B) \rightarrow (\forall B^*. A B) : \star$$
$$\vdash_{\emptyset} \lambda C^*. C : \star \rightarrow \star$$

$$\not\vdash_{\emptyset} \lambda H^{\forall B^*. (\lambda C^*. C) B}. H : (\forall B^*. (\lambda C^*. C) B) \rightarrow (\forall B^*. (\lambda C^*. C) B)$$

where $\star := \text{Prop}$

Toward a Pedagogical Calculus of Constructions

Summary of necessary constraints

$$\frac{\Gamma \vdash A : s_1 \quad \Gamma, x : A \vdash B : s_2}{\Gamma \vdash \forall x^A. B : s_2} \text{ (prod)}$$

(s_1, s_2)	λ^2	λ^ω
(Prop, Prop)	✓	✓
(Type, Prop)	✗	✗
(Prop, Type)		
(Type, Type)		✓

- ▶ ✓ instance of the rule does not produce empty type
- ▶ ✗ instance of the rule can produce empty types

Toward a Pedagogical Calculus of Constructions

Summary of necessary constraints

$$\frac{\Gamma \vdash A : s_1 \quad \Gamma, x : A \vdash B : s_2}{\Gamma \vdash \forall x^A. B : s_2} \text{ (prod)}$$

(s_1, s_2)	λ^2	λ^ω	λC
(Prop, Prop)	✓	✓	✗
(Type, Prop)	✗	✗	✗
(Prop, Type)			✓
(Type, Type)		✓	✓

- ▶ ✓ instance of the rule does not produce empty type
- ▶ ✗ instance of the rule can produce empty types

Toward a Pedagogical Calculus of Constructions

Summary of necessary constraints

$$\frac{\Gamma \vdash v : A : s_1 \quad \Gamma \vdash u : \forall x^A. B : s_2}{\Gamma \vdash u \ v : B[x \leftarrow v]} \text{ (app)}$$

(s_1, s_2)	λ^2	λ^ω
(Prop, Prop)	✓	✓
(Type, Prop)	✓	✗
(Prop, Type)		
(Type, Type)		✗

- ▶ ✓ instance of the rule does not produce empty type
- ▶ ✗ instance of the rule can produce empty types

Toward a Pedagogical Calculus of Constructions

Summary of necessary constraints

$$\frac{\Gamma \vdash v : A : s_1 \quad \Gamma \vdash u : \forall x^A. B : s_2}{\Gamma \vdash u \ v : B[x \leftarrow v]} \text{ (app)}$$

(s_1, s_2)	λ^2	λ^ω	λC
(Prop, Prop)	✓	✓	✗
(Type, Prop)	✓	✗	✗
(Prop, Type)			✗
(Type, Type)		✗	✗

- ▶ ✓ instance of the rule does not produce empty type
- ▶ ✗ instance of the rule can produce empty types

Toward a Pedagogical Calculus of Constructions

Pedagogical Calculus of Constructions?

$$\frac{}{\emptyset \text{wf}_{\sigma}^{\text{Ce}}} \text{(env}_1\text{)}$$

$$\frac{\Gamma \vdash_{\sigma}^{\text{Ce}} A : \kappa \quad \vdash_{\emptyset}^{\text{Ce}} a : A' \quad \sigma(A) \rightsquigarrow^* A' \quad x \notin \text{dom}(\Gamma)}{\Gamma, x : A \text{wf}_{\sigma::(x \mapsto a)}^{\text{Ce}}} \text{(env}_2\text{)}$$

$$\frac{\Gamma \text{wf}_{\sigma}^{\text{Ce}}}{\Gamma \vdash_{\sigma}^{\text{Ce}} \circ : \top : \text{Prop} : \text{Type}} \text{(ax)}$$

$$\frac{\Gamma, x : A, \Gamma' \text{wf}_{\sigma}^{\text{Ce}}}{\Gamma, x : A, \Gamma' \vdash_{\sigma}^{\text{Ce}} x : A} \text{(var)}$$

$$\frac{\Gamma, x : A \vdash_{\sigma::(x \mapsto a)}^{\text{Ce}} u : B : \kappa}{\Gamma \vdash_{\sigma}^{\text{Ce}} \lambda x^A. u : \forall x^A. B} \text{(abs)}$$

Toward a Pedagogical Calculus of Constructions

Pedagogical Calculus of Constructions?

$$\frac{\Gamma \vdash_{\sigma}^{\text{Ce}} u : \forall x^A. B : \text{Prop} \quad \Gamma \vdash_{\sigma}^{\text{Ce}} v : A \quad \sigma \text{ mot}^{\text{Prop}}(B[x \leftarrow v])}{\Gamma \vdash_{\sigma}^{\text{Ce}} u v : B[x \leftarrow v]} \text{ (app}_{\star}\text{)}$$

$$\frac{\Gamma \vdash_{\sigma}^{\text{Ce}} u : \forall x^A. B : \text{Type} \quad \Gamma \vdash_{\sigma}^{\text{Ce}} v : A \quad \sigma \text{ mot}^{B[x \leftarrow v]}(u v)}{\Gamma \vdash_{\sigma}^{\text{Ce}} u v : B[x \leftarrow v]} \text{ (app}_{\square}\text{)}$$

$$\frac{\Gamma, x : A \vdash_{\sigma :: (x \mapsto a)}^{\text{Ce}} B : \text{Prop} \quad \sigma \text{ mot}^{\text{Prop}}(\forall x^A. B)}{\Gamma \vdash_{\sigma}^{\text{Ce}} \forall x^A. B : \text{Prop}} \text{ (prod}_{\star}\text{)}$$

$$\frac{\Gamma, x : A \vdash_{\sigma :: (x \mapsto a)}^{\text{Ce}} B : \text{Type}}{\Gamma \vdash_{\sigma}^{\text{Ce}} \forall x^A. B : \text{Type}} \text{ (prod}_{\square}\text{)}$$

$$\frac{\Gamma \vdash_{\sigma}^{\text{Ce}} u : A \quad A \simeq A' \quad \Gamma \vdash_{\sigma}^{\text{Ce}} A' : \kappa}{\Gamma \vdash_{\sigma}^{\text{Ce}} u : A'} \text{ (conv)}$$

Conclusion and further work

Contributions

- ▶ formal definition of pedagogical sub-system of CC
- ▶ examples of this definition (λ_e^2 , λ_t^2 , λ_p^2 , λ_e^ω)
- ▶ introduction of explicit exemplifications
- ▶ study of type checking pedagogical calculi
- ▶ motivated conjectures : pedagogical higher-order and Calculus of Constructions

Further work

- ▶ show the conjectures
- ▶ study machine implementation
- ▶ extend to stronger systems
- ▶ formal verification of Griss' negationless mathematics
- ▶ inconsistency and pedagogy...